

## Références

- [1] Tandon R, Keshavan MS, Nasrallah HA. Schizophrenia, “just the facts”: What we know in 2008. Part 1: Overview. *Schizophr Res* 2008;100:4–19.
- [2] Tandon R, Keshavan MS, Nasrallah HA. Schizophrenia, “just the facts” what we know in 2008. Part 2: Epidemiology and etiology. *Schizophr Res* 2008;102:1–18.
- [3] MacDonald AW, Schulz SC. What we know: Findings that every theory of schizophrenia should explain. *Schizophr Bull* 2009;35:493–508.
- [4] Weinberger DR. Implications of normal brain development for the pathogenesis of schizophrenia. *Arch Gen Psychiatry* 1987;44:660–9.
- [5] Fatemi SH, Folsom TD. The neurodevelopmental hypothesis of schizophrenia, revisited. *Schizophr Bull* 2009;35:528–48.
- [6] Lecomte T, Leclerc C, Corbiere M, Wykes T, Wallace CJ, Spidel A. Group cognitive behavior therapy or social skills training for individuals with a recent onset of psychosis? Results of a randomized controlled trial. *J Nerv Ment Dis* 2008;196: 866–75.
- [7] Sells DJ, Stayner DA, Davidson L. Recovering the self in schizophrenia: An integrative review of qualitative studies. *Psychiatr Q* 2004;75:87–97.
- [8] Favrod J, Scheder D. Se rétablir de la schizophrénie : un modèle d'intervention. *Rev Med Suisse Romande* 2004;124:205–8.
- [9] Andresen R, Oades L, Caputi P. The experience of recovery from schizophrenia: Towards an empirically validated stage model. *Aust N Z J Psychiatry* 2003;37:586–94.
- [10] Liberman R, Kopelowicz A, Ventura J, Gutkind D. Operational criteria and factors related to recovery from schizophrenia. *International Review of Psychiatry* 2003;14: 256–72.
- [11] Tempier R, Favrod J. Réadaptation psychiatrique en milieu francophone : pratiques actuelles, défis futurs. *Can J Psychiatry* 2002;47:621–7.
- [12] Bonsack C, Adam L, Haefliger T, Besson J, Conus P. Difficult-to-engage patients: A specific target for time-limited assertive outreach in a swiss setting. *Can J Psychiatry* 2005;50:845–50.
- [13] Bonsack C, Haefliger T, Cordier S, Conus P. Accès aux soins et maintien dans la communauté des personnes difficiles à engager dans un traitement psychiatrique. *Rev Med Suisse Romande* 2004;124:225–9.
- [14] Kanter J. Clinical case management: Definition, principles, components. *Hosp Community Psychiatry* 1989;40:361–8.
- [15] DiClemente CC, Prochaska JO. Self-change and therapy change of smoking behavior: A comparison of processes of change in cessation and maintenance. *Addict Behav* 1982;7:133–42.
- [16] Prochaska JO, DiClemente CC. Stages and processes of self-change of smoking: Toward an integrative model of change. *J Consult Clin Psychol* 1983;51:390–5.
- [17] Prochaska JO, DiClemente CC. Self change processes, self efficacy and decisional balance across five stages of smoking cessation. *Prog Clin Biol Res* 1984;156:131–40.
- [18] Linszen DH, Dingemans PM, Lenior ME. Cannabis abuse and the course of recent-onset schizophrenic disorders. *Arch Gen Psychiatry* 1994;51:273–9.
- [19] Carpenter WT Jr, Strauss JS, Muleh S. Are there pathognomonic symptoms in schizophrenia? An empiric investigation of schneider's first-rank symptoms. *Arch Gen Psychiatry* 1973;28:847–52.
- [20] David AS. Insight and psychosis. *Br J Psychiatry* 1990;156:798–808.

- [21] Aleman A, de Haan EH, Kahn RS. Insight and neurocognitive function in schizophrenia. *J Neuropsychiatry Clin Neurosci* 2002;14:241–2.
- [22] Subotnik KL, Nuechterlein KH, Irzhevsky V, Kitchen CM, Woo SM, Mintz J. Is unawareness of psychotic disorder a neurocognitive or psychological defensiveness problem? *Schizophr Res* 2005;75:147–57.
- [23] Cooke MA, Peters ER, Kuipers E, Kumari V. Disease, deficit or denial? Models of poor insight in psychosis. *Acta Psychiatr Scand* 2005;112:4–17.
- [24] Green MF, Kern RS, Robertson MJ, Sergi MJ, Kee KS. Relevance of neurocognitive deficits for functional outcome in schizophrénie. In: Sharma T, Harvey PD, editors. *Cognition in schizophrenia*. New York: Oxford University Press; 2000. p. 178–92.
- [25] Startup M. Awareness of own and others' schizophrenic illness. *Schizophr Res* 1997;26:203–11.
- [26] Lysaker PH, Lancaster RS, Davis LW, Clements CA. Patterns of neurocognitive deficits and unawareness of illness in schizophrenia. *J Nerv Ment Dis* 2003;191:38–44.
- [27] Cunningham K, Wolbert R, Brockmeier MB. Moving beyond the illness: Factors contributing to gaining and maintaining employment. *Am J Community Psychol* 2000;28:481–94.
- [28] Conley RR, Ascher-Svanum H, Zhu B, Faries DE, Kinon BJ. The burden of depressive symptoms in the long-term treatment of patients with schizophrenia. *Schizophr Res* 2007;90:186–97.
- [29] Potvin S, Sepehry AA, Stip E. Meta-analysis of depressive symptoms in dual-diagnosis schizophrenia. *Aust N Z J Psychiatry* 2007;41:792–9.
- [30] Rocca P, Bellino S, Calvarese P, Marchiaro L, Patria L, Rasetti R, Bogetto F. Depressive and negative symptoms in schizophrenia: Different effects on clinical features. *Compr Psychiatry* 2005;46:304–10.
- [31] Zisook S, McAdams LA, Kuck J, Harris MJ, Bailey A, Patterson TL, et al. Depressive symptoms in schizophrenia. *Am J Psychiatry* 1999;156:1736–43.
- [32] Brebion G, Smith MJ, Amador X, Malaspina D, Gorman JM. Clinical correlates of memory in schizophrenia: Differential links between depression, positive and negative symptoms, and two types of memory impairment. *Am J Psychiatry* 1997;154:1538–43.
- [33] Haw C, Hawton K, Sutton L, Sinclair J, Deeks J. Schizophrenia and deliberate self-harm: A systematic review of risk factors. *Suicide Life Threat Behav* 2005;35:50–62.
- [34] Palmer BA, Pankratz VS, Bostwick JM. The lifetime risk of suicide in schizophrenia: A reexamination. *Arch Gen Psychiatry* 2005;62:247–53.
- [35] Addington D, Addington J, Atkinson M. A psychometric comparison of the calgary depression scale for schizophrenia and the hamilton depression rating scale. *Schizophr Res* 1996;19:205–12.
- [36] Kim SW, Kim SJ, Yoon BH, Kim JM, Shin IS, Hwang MY, Yoon JS. Diagnostic validity of assessment scales for depression in patients with schizophrenia. *Psychiatry Res* 2006;144:57–63.
- [37] Muller MJ, Muller KM, Fellgiebel A. Detection of depression in acute schizophrenia : Sensitivity and specificity of 2 standard observer rating scales. *Can J Psychiatry* 2006;51:387–92.
- [38] Lançon C, Auquier P, Reine G, Bernard D, Toumi M. Study of the concurrent validity of the calgary depression scale for schizophrenics (cdss). *J Affect Disord* 2000;58:107–15.
- [39] Lançon C, Auquier P, Reine G, Toumi M, Addington D. Evaluation of depression in schizophrenia: Psychometric properties of a french version of the calgary depression scale. *Psychiatry Res* 1999;89:123–32.
- [40] Reine G, Bernard D, Auquier P, Le Fur B, Lancon C. Propriétés psychométriques de la version française de l'échelle de dépression de calgary pour schizophrénies (cdss). *Encephale* 2000;26:52–61.

- [41] Cuijpers P, Munoz RF, Clarke GN, Lewinsohn PM. Psychoeducational treatment and prevention of depression: The "coping with depression" course thirty years later. *Clin Psychol Rev* 2009;29(5):449–58.
- [42] Dimeo F, Bauer M, Varahram I, Proest G, Halter U. Benefits from aerobic exercise in patients with major depression: A pilot study. *Br J Sports Med* 2001;35:114–7.
- [43] Knubben K, Reischies FM, Adli M, Schlattmann P, Bauer M, Dimeo F. A randomised, controlled study on the effects of a short-term endurance training programme in patients with major depression. *Br J Sports Med* 2007;41:29–33.
- [44] Walsh E, Buchanan A, Fahy T. Violence and schizophrenia: Examining the evidence. *Br J Psychiatry* 2002;180:490–5.
- [45] Swanson JW, Holzer CE 3rd, Ganju VK, Jono RT. Violence and psychiatric disorder in the community: Evidence from the epidemiologic catchment area surveys. *Hosp Community Psychiatry* 1990;41:761–70.
- [46] Tiihonen J, Isohanni M, Rasanen P, Koiranen M, Moring J. Specific major mental disorders and criminality: A 26-year prospective study of the 1966 northern finland birth cohort. *Am J Psychiatry* 1997;154:840–5.
- [47] Brennan PA, Mednick SA, Hodgins S. Major mental disorders and criminal violence in a danish birth cohort. *Arch Gen Psychiatry* 2000;57:494–500.
- [48] Wallace C, Mullen P, Burgess P. Serious criminal offending and mental disorder. Case linkage study. *Br J Psychiatry* 1998;172:477–84.
- [49] Wootton L, Buchanan A, Leese M, Tyrer P, Burns T, Creed F, et al. Violence in psychosis : Estimating the predictive validity of readily accessible clinical information in a community sample. *Schizophr Res* 2008;101(1–3):176–84.
- [50] Regier DA, Farmer ME, Rae DS, Locke BZ, Keith SJ, Judd LL, Goodwin FK. Comorbidity of mental disorders with alcohol and other drug abuse. Results from the epidemiologic catchment area (eca) study. *JAMA* 1990;264:2511–8.
- [51] Tucker P. Substance misuse and early psychosis. *Australas Psychiatry* 2009;17(4):291–4.
- [52] Lambert M, Conus P, Lubman DI, Wade D, Yuen H, Moritz S, et al. The impact of substance use disorders on clinical outcome in 643 patients with first-episode psychosis. *Acta Psychiatr Scand* 2005;112:141–8.
- [53] Rathbone J, Variend H, Mehta H. Cannabis and schizophrenia. *Cochrane Database Syst Rev* 2008 : CD004837.
- [54] Zammit S, Moore TH, Lingford-Hughes A, Barnes TR, Jones PB, Burke M, Lewis G. Effects of cannabis use on outcomes of psychotic disorders: Systematic review. *Br J Psychiatry* 2008;193:357–63.
- [55] Leucht S, Cipriani A, Spineli L, Mavridis D, Orey D, Richter F, et al. Comparative efficacy and tolerability of 15 antipsychotic drugs in schizophrenia: A multiple-treatments meta-analysis. *Lancet* 2013;382:951–62.
- [56] Leucht S, Tardy M, Komossa K, Heres S, Kissling W, Salanti G, Davis JM. Antipsychotic drugs versus placebo for relapse prevention in schizophrenia: A systematic review and meta-analysis. *Lancet* 2012;379:2063–71.
- [57] Harrow M, Jobe TH. Does long-term treatment of schizophrenia with antipsychotic medications facilitate recovery? *Schizophr Bull* 2013;39:962–5.
- [58] Andreasen NC, Carpenter WT Jr, Kane JM, Lasser RA, Marder SR, Weinberger DR. Remission in schizophrenia: Proposed criteria and rationale for consensus. *Am J Psychiatry* 2005;162:441–9.
- [59] Viguera AC, Baldessarini RJ, Hegarty JD, van Kammen DP, Tohen M. Clinical risk following abrupt and gradual withdrawal of maintenance neuroleptic treatment. *Arch Gen Psychiatry* 1997;54:49–55.
- [60] Moncrieff J. Does antipsychotic withdrawal provoke psychosis? Review of the literature on rapid onset psychosis (supersensitivity psychosis) and withdrawal-related relapse. *Acta Psychiatr Scand* 2006;114:3–13.

- [61] Harrow M, Jobe TH, Faull RN. Do all schizophrenia patients need antipsychotic treatment continuously throughout their lifetime? A 20-year longitudinal study. *Psychol Med* 2012;42:2145–55.
- [62] Byerly MJ, Nakonezny PA, Lescouffair E. Antipsychotic medication adherence in schizophrenia. *Psychiatr Clin North Am* 2007;30:437–52.
- [63] Karow A, Czekalla J, Dittmann RW, Schacht A, Wagner T, Lambert M, et al. Association of subjective well-being, symptoms, and side effects with compliance after 12 months of treatment in schizophrenia. *J Clin Psychiatry* 2007;68:75–80.
- [64] Moritz S, Peters MJ, Karow A, Deljkovic A, Tonn P, Naber D. Cure or curse? Ambivalent attitudes towards neuroleptic medication in schizophrenia and non-schizophrenia patients. *Mental Illness* 2009;1:e2.
- [65] Moritz S, Favrod J, Andreou C, Morrison AP, Bohn F, Veckenstedt R, et al. Beyond the usual suspects: Positive attitudes towards positive symptoms is associated with medication noncompliance in psychosis. *Schizophr Bull* 2013;39:917–22.
- [66] McGorry PD, Chanan A, McCarthy E, Van Riel R, McKenzie D, Singh BS. Posttraumatic stress disorder following recent-onset psychosis. An unrecognized postpsychotic syndrome. *J Nerv Ment Dis* 1991;179:253–8.
- [67] Morrison AP, Frame L, Larkin W. Relationships between trauma and psychosis: A review and integration. *Br J Clin Psychol* 2003;42:331–53.
- [68] Tarrier N, Khan S, Cater J, Picken A. The subjective consequences of suffering a first episode psychosis: Trauma and suicide behaviour. *Soc Psychiatry Psychiatr Epidemiol* 2007;42:29–35.
- [69] Mizrahi R, Rusjan P, Agid O, Graff A, Mamo DC, Zipursky RB, Kapur S. Adverse subjective experience with antipsychotics and its relationship to striatal and extrastriatal d2 receptors: A pet study in schizophrenia. *Am J Psychiatry* 2007;164:630–7.
- [70] Marder SR. Subjective experiences on antipsychotic medications: Synthesis and conclusions. *Acta Psychiatr Scand Suppl* 2005;43–6.
- [71] Voruganti LN, Awad AG. Subjective and behavioural consequences of striatal dopamine depletion in schizophrenia—findings from an in vivo spect study. *Schizophr Res* 2006;88:179–86.
- [72] Lecomte T, Wallace CJ, Perreault M, Caron J. Consumers' goals in psychiatric rehabilitation and their concordance with existing services. *Psychiatr Serv* 2005;56:209–11.
- [73] Overall JE. The brief psychiatric rating scale (bprs): Recent developments in ascertainment and scaling. *Psychopharmacol Bull* 1988;24:97–9.
- [74] Lukoff D, Liberman RP, Nuechterlein KH. Symptom monitoring in the rehabilitation of schizophrenic patients. *Schizophr Bull* 1986;12:578–602.
- [75] Andreasen NC. The scale for the assessment of negative symptoms (SANS): Conceptual and theoretical foundations. *Br J Psychiatry Suppl* 1989;49–58.
- [76] Favrod J, Rexhaj S, Ferrari P, Bardy S, Hayoz C, Morandi S, et al. French version validation of the psychotic symptom rating scales (PSYRATS) for outpatients with persistent psychotic symptoms. *BMC Psychiatry* 2012;12:161.
- [77] Haddock G, McCarron J, Tarrier N, Faragher EB. Scales to measure dimensions of hallucinations and delusions: The psychotic symptom rating scales (PSYRATS). *Psychol Med* 1999;29:879–89.
- [78] Favrod J, Grasset F, Spreng S, Grossenbacher B, Hodé Y. Benevolent voices are not so kind: The functional significance of auditory hallucinations. *Psychopathology* 2004;37:304–8.
- [79] Favrod J, Pomini V, Grasset F. Thérapie cognitive et comportementale pour les hallucinations auditives résistantes au traitement neuroleptique. *Rev Med Suisse Romande* 2004;124:213–6.
- [80] Favrod J, Zimmermann G, Raffard S, Pomini V, Khazaal Y. The beck cognitive insight scale in outpatients with psychotic disorders: Further evidence from a french-speaking sample. *Can J Psychiatry* 2008;53:783–7.

- 
- [81] Aubin G, Stip E, Gelinias I, Rainville C, Chapparo C. Daily activities, cognition and community functioning in persons with schizophrenia. *Schizophr Res* 2009;107:313–8.
  - [82] Cyr M, Toupin J, Lesage AD, Valiquette CA. Assessment of independent living skills for psychotic patients. Further validity and reliability. *J Nerv Ment Dis* 1994;182:91–7.
  - [83] Wallace CJ, Liberman RP, Tauber R, Wallace J. The independent living skills survey: A comprehensive measure of the community functioning of severely and persistently mentally ill individuals. *Schizophr Bull* 2000;26:631–58.
  - [84] Favrod J, McQuillan A, Pomini V, Ferrero FP. Training interpersonal problem-solving skills in french-speaking switzerland. *International Review of Psychiatry* 1998;10:30–4.
  - [85] Mohr S, Simon A, Favrod J, Fokianos C, Ferrero F. Validation de la version française du profil des compétences de vie (life skills profile) chez des personnes atteintes de schizophrénie. *Encephale* 2004;30:343–51.
  - [86] Pomini V, Golay P, Reymond C. L'évaluation des difficultés et des besoins des patients psychiatriques. *L'Information Psychiatrique* 2008;84:895–902.
  - [87] Bonsack C, Montagnin Y, Favrod J, Gibellini S, Conus P. Une intervention motivationnelle pour les consommateurs de cannabis souffrant de psychose. *Encephale* 2007;33:819–26.
  - [88] Lacroix A, Assal JP. L'éducation thérapeutique des patients. Paris: Maloine; 2003.
  - [89] D'Ivernoy JF, Gagnayre R. Apprendre à éduquer le patient : Approche pédagogique. Paris: Maloine; 2004.
  - [90] Chéreau I, Llorca PM, Petit M. L'intérêt des programmes psychoéducatifs. *Santé Mentale* 2008;133:48–9.
  - [91] Goldman CR. Toward a definition of psychoeducation. *Hosp Community Psychiatry* 1988;39:666–8.
  - [92] Carling PJ, Palmer H. Dans l'œil du cyclone : des assises à actualiser en santé communautaire. In: Emard R, Aubry T, editors. *Le suivi communautaire en santé mentale : une invitation à bâtir sa vie*. Ottawa: Les Presses de l'Université d'Ottawa; 2004. p. 279–99.
  - [93] Mueser KT, Bond GR, Drake RE, Resnick SG. Models of community care for severe mental illness: A review of research on case management. *Schizophr Bull* 1998;24:37–74.
  - [94] Lamb HR, Mackota C. Vocational services in a community mental health program. *Hosp Community Psychiatry* 1968;19:315–8.
  - [95] Liberman RP, McCann MJ, Wallace CJ. Generalization of behaviour therapy with psychotics. *Br J Psychiatry* 1976;129:490–6.
  - [96] Anthony WA, Liberman RP. The practice of psychiatric rehabilitation: Historical, conceptual, and research base. *Schizophr Bull* 1986;12:542–59.
  - [97] Anderson CM, Hogarty GE, Reiss DJ. Family treatment of adult schizophrenic patients : A psycho-educational approach. *Schizophr Bull* 1980;6:490–505.
  - [98] Gélinas D. L'embauche d'usagers à titre de pourvoyeurs de services de santé mentale. *Le Partenaire* 2006;14:9–41.
  - [99] Deegan PE. Recovery: The lived experience of rehabilitation. *Psychosocial Rehabilitation Journal* 1988;11:11–9.
  - [100] Deegan PE. Recovery as a journey of heart. *Psychiatric Rehabilitation Journal* 1996;19:91–7.
  - [101] Deegan PE. Recovery and empowerment for people with psychiatric disabilities. *Soc Work Health Care* 1997;25:11–24.
  - [102] Mead S, Copeland ME. What recovery means to us: Consumer's perspectives. *Community Ment Health J* 2000;36:315–28.
  - [103] Deegan PE. Le rétablissement comme un processus de transformation. *Santé Mentale* 2006;106:68–76.
  - [104] Rogers ES, Chamberlin J, Ellison ML, Crean T. A consumer-constructed scale to measure empowerment among users of mental health services. *Psychiatr Serv* 1997;48:1042–7.

- [105] Cook JA, Copeland ME, Hamilton MM, Jonikas JA, Razzano LA, Floyd CB, et al. Initial outcomes of a mental illness self-management program based on wellness recovery action planning. *Psychiatr Serv* 2009;60:246–9.
- [106] Carling PJ. Housing and supports for persons with mental illness: Emerging approaches to research and practice. *Hosp Community Psychiatry* 1993;44:439–49.
- [107] Rapp CA. The strenght model. New-York: Oxford University Press; 1998.
- [108] LeRoy S. Recovery from psychiatric disability: Implications for rehabilitation counseling education. *Rehabilitation Education* 2001;15:167–75.
- [109] Provencher HL. Le paradigme du rétablissement: 1. Une expérience globale de santé. *Le Partenaire* 2007;15.
- [110] McCann TV, Clark E. Advancing self-determination with young adults who have schizophrenia. *J Psychiatr Ment Health Nurs* 2004;11:12–20.
- [111] Provencher HL. Le rétablissement : un nouveau modèle en santé mentale. *Santé Mentale* 2006;106:59–65.
- [112] Askey R, Gamble C, Gray R. Family work in first-onset psychosis: A literature review. *J Psychiatr Ment Health Nurs* 2007;14:356–65.
- [113] Valimaki M, Leino-Kilpi H. Preconditions for and consequences of self-determination: The psychiatric patient's point of view. *J Adv Nurs* 1998;27:204–12.
- [114] Yeo M, Williams JR, Hooper W. Incorporating ethics in priority setting : A case study of a regional health board in canada. *Health Care Anal* 1999;7:177–94.
- [115] Henry L, Edwards J, Jackson H, Hulbert C, MCGorry P. Cognitively oriented psychotherapy for first episod psychosis (cope): A practioner's manual. Melbourne: Early Psychosis Prevention and Intervention Center; 2002.
- [116] EPPIC. Psychoeducation in early psychosis: A manual. Melbourne: Early Psychosis Prevention and Intervention Center; 2001.
- [117] Corrigan PW, Wassel A. Understanding and influencing the stigma of mental illness. *J Psychosoc Nurs Ment Health Serv* 2008;46:42–8.
- [118] Tait L, Birchwood M, Trower P. Predicting engagement with services for psychosis: Insight, symptoms and recovery style. *Br J Psychiatry* 2003;182:123–8.
- [119] Thompson KN, MCGorry PD, Harrigan SM. Recovery style and outcome in first-episode psychosis. *Schizophr Res* 2003;62:31–6.
- [120] Tait L, Birchwood M, Trower P. Adapting to the challenge of psychosis: Personal resilience and the use of sealing-over (avoidant) coping strategies. *Br J Psychiatry* 2004;185:410–5.
- [121] Herman NJ. Return to sender. *Journal of Contemporary Ethnography* 1993;22: 295–330.
- [122] Gleeson J, Jackson HJ, Stavelly H, Burnett P. Family intervention in early psychosis. In: MCGorry P, Jackson H, editors. *The recognition and management of early psychosis: A preventive approach*. Cambridge: Cambridge University Press; 1999. p. 376–406.
- [123] Hartog G, Dufort F. Les défis de l'intervention dans un contexte multiethnique. In: Dufort F, Guay J, editors. *Agir au cœur des communautés : la psychologie communautaire et le changement social*. Québec: Les Presses de l'Université Laval; 2002.
- [124] Cohen-Emerique M. [the intercultural approach in the support process]. *Sante Ment Que* 1993;18:71–91.
- [125] Hooley JM. Expressed emotion and locus of control. *J Nerv Ment Dis* 1998;186:374–8.
- [126] Hooley JM. Expressed emotion and relapse of psychopathology. *Annu Rev Clin Psychol* 2007;3:329–52.
- [127] Pekkala E, Merinder L. Psychoeducation for schizophrenia. *Cochrane Database Syst Rev* 2002 : CD002831.
- [128] Magliano L, Fiorillo A, Fadden G, Gair F, Economou M, Kallert T, et al. Effectiveness of a psychoeducational intervention for families of patients with schizophrenia: Preliminary results of a study funded by the european commission. *World Psychiatry* 2005;4:45–9.



- 
- [129] Sin J, Moone N, Newell J. Developing services for the carers of young adults with early-onset psychosis - implementing evidence-based practice on psycho-educational family intervention. *J Psychiatr Ment Health Nurs* 2007;14:282-90.
  - [130] Cottraux J. Les thérapies comportementales et cognitives. Paris: Masson; 1998.
  - [131] Young JE, Klosko JS, Weishaar ME. La thérapie des schémas : approche cognitive des troubles de la personnalité. Bruxelles: De Boeck Université; 2006.
  - [132] Liberman RP. Éducation au contrôle des symptômes. Charleroi: Socrates Editions Promarex; 1995.
  - [133] Hatam S. Directives anticipées : prévoir une incapacité de discernement, rédiger des directives anticipées. Genève: Pro Mente Sana; 2007.
  - [134] Khazaal Y. Repérer les prodromes de la manie et prévenir la récurrence. *Santé Mentale* 2007;127:56-60.
  - [135] Khazaal Y, Richard C, Matthieu-Darekar S, Quement B, Kramer U, Preisig M. Advance directives in bipolar disorder, a cognitive behavioural conceptualization. *Int J Law Psychiatry* 2008;31:1-8.
  - [136] Rubio F. Conduites d'entretiens d'investigations, motivationnels, psychothérapeutiques. Lausanne : Département Universitaire de Psychiatrie Adulte, CHUV 2000;.
  - [137] Jackson C, Knott C, Skeate A, Birchwood M. The trauma of first episode psychosis : The role of cognitive mediation. *Aust N Z J Psychiatry* 2004;38:327-33.
  - [138] Steil R, Ehlers A. Dysfunctional meaning of posttraumatic intrusions in chronic ptsd. *Behav Res Ther* 2000;38:537-58.
  - [139] Mueser KT, Rosenberg SD, Goodman LA, Trumbetta SL. Trauma, ptsd, and the course of severe mental illness: An interactive model. *Schizophr Res* 2002;53:123-43.
  - [140] Mueser KT, Rosenberg SD. Treating the trauma of first episode psychosis: A ptsd perspective. *J Ment Health* 2003;12:103-8.
  - [141] Rademaker AR, Vermetten E, Geuze E, Mulwijk A, Kleber RJ. Self-reported early trauma as a predictor of adult personality: A study in a military sample. *J Clin Psychol* 2008;64:863-75.
  - [142] Strauss JS. Hallucinations and delusions as points on continua function. Rating scale evidence. *Arch Gen Psychiatry* 1969;21:581-6.
  - [143] Verdoux H, van Os J. Psychotic symptoms in non-clinical populations and the continuum of psychosis. *Schizophr Res* 2002;54:59-65.
  - [144] Peters E, Joseph S, Day S, Garety P. Measuring delusional ideation: The 21-item peters et al. Delusions inventory (PDI). *Schizophr Bull* 2004;30:1005-22.
  - [145] Johns LC. Hallucinations in the general population. *Curr Psychiatry Rep* 2005;7:162-7.
  - [146] Johns LC, van Os J. The continuity of psychotic experiences in the general population. *Clin Psychol Rev* 2001;21:1125-41.
  - [147] Freeman D, Pugh K, Garety P. Jumping to conclusions and paranoid ideation in the general population. *Schizophr Res* 2008;102:254-60.
  - [148] Warman DM, Lysaker PH, Martin JM, Davis L, Haudenschild SL. Jumping to conclusions and the continuum of delusional beliefs. *Behav Res Ther* 2007;45:1255-69.
  - [149] Milton F, Patwa VK, Hafner RJ. Confrontation vs. Belief modification in persistently deluded patients. *Br J Med Psychol* 1978;51:127-30.
  - [150] Watts FN, Powell GE, Austin SV. The modification of abnormal beliefs. *Br J Med Psychol* 1973;46:359-63.
  - [151] Moller P, Husby R. The initial prodrome in schizophrenia : Searching for naturalistic core dimensions of experience and behavior. *Schizophr Bull* 2000;26:217-32.
  - [152] Stone JM, Pilowsky LS. Antipsychotic drug action: Targets for drug discovery with neurochemical imaging. *Expert Rev Neurother* 2006;6:57-64.
  - [153] Conley RR, Kelly DL. Management of treatment resistance in schizophrenia. *Biol Psychiatry* 2001;50:898-911.
  - [154] Miller LJ. Qualitative changes in hallucinations. *Am J Psychiatry* 1996;153:265-7.

- [155] Mizrahi R, Kiang M, Mamo DC, Arenovich T, Bagby RM, Zipursky RB, Kapur S. The selective effect of antipsychotics on the different dimensions of the experience of psychosis in schizophrenia spectrum disorders. *Schizophr Res* 2006;88:111–8.
- [156] Blakemore SJ. Deluding the motor system. *Conscious Cogn* 2003;12:647–55.
- [157] Franck N, Rouby P, Daprati E, Dalery J, Marie-Cardine M, Georgieff N. Confusion between silent and overt reading in schizophrenia. *Schizophr Res* 2000;41:357–64.
- [158] Franck N, Farrer C, Georgieff N, Marie-Cardine M, Dalery J, d'Amato T, Jeannerod M. Defective recognition of one's own actions in patients with schizophrenia. *Am J Psychiatry* 2001;158:454–9.
- [159] Lecomte T, Spidel A, Leclerc C, MacEwan GW, Greaves C, Bentall RP. Predictors and profiles of treatment non-adherence and engagement in services problems in early psychosis. *Schizophr Res* 2008;102:295–302.
- [160] Maher BA. Delusional thinking and perceptual disorder. *J Individ Psychol* 1974;30:98–113.
- [161] Bell V, Halligan PW, Ellis HD. The cardiff anomalous perceptions scale (caps): A new validated measure of anomalous perceptual experience. *Schizophr Bull* 2006;32:366–77.
- [162] Lecomte T, Liberman RP, Wallace CJ. Identifying and using reinforcers to enhance the treatment of persons with serious mental illness. *Psychiatr Serv* 2000;51:1312–4.
- [163] Pelak VS, Liu GT. Visual hallucinations. *Curr Treat Options Neurol* 2004;6:75–83.
- [164] Baethge C. Grief hallucinations: True or pseudo? Serious or not? An inquiry into psychopathological and clinical features of a common phenomenon. *Psychopathology* 2002;35:296–302.
- [165] Tanriverdi N, Sayilgan MA, Ozcurumez G. Musical hallucinations associated with abruptly developed bilateral loss of hearing. *Acta Psychiatr Scand* 2001;103:153–5.
- [166] Baba A, Hamada H. Musical hallucinations in schizophrenia. *Psychopathology* 1999;32:242–51.
- [167] Baba A, Hamada H, Kocha H. Musical hallucinations in schizophrenia. 2. Relations with verbal hallucinations. *Psychopathology* 2003;36:104–10.
- [168] Saba PR, Keshavan MS. Musical hallucinations and musical imagery: Prevalence and phenomenology in schizophrenic inpatients. *Psychopathology* 1997;30:185–90.
- [169] Eisen JL, Phillips KA, Baer L, Beer DA, Atala KD, Rasmussen SA. The brown assessment of beliefs scale: Reliability and validity. *Am J Psychiatry* 1998;155:102–8.
- [170] Daprati E, Franck N, Georgieff N, Proust J, Pacherie E, Dalery J, Jeannerod M. Looking for the agent: An investigation into consciousness of action and self-consciousness in schizophrenic patients. *Cognition* 1997;65:71–86.
- [171] Favrod J, Vianin P, Pomini V. Reconnaissance de la source et symptômes psychotiques : implications pour la clinique. *Revue Francophone de Clinique Comportementale et Cognitive* 2008;13:8–17.
- [172] Vinogradov S, Willis-Shore J, Poole JH, Marten E, Ober BA, Shenaut GK. Clinical and neurocognitive aspects of source monitoring errors in schizophrenia. *Am J Psychiatry* 1997;154:1530–7.
- [173] Keefe RS, Arnold MC, Bayen UJ, McEvoy JP, Wilson WH. Source-monitoring deficits for self-generated stimuli in schizophrenia: Multinomial modeling of data from three sources. *Schizophr Res* 2002;57:51–67.
- [174] Morrison AP, Haddock G. Cognitive factors in source monitoring and auditory hallucinations. *Psychol Med* 1997;27:669–79.
- [175] Allen PP, Johns LC, Fu CH, Broome MR, Vythelingum GN, McGuire PK. Misattribution of external speech in patients with hallucinations and delusions. *Schizophr Res* 2004;69:277–87.
- [176] Bentall RP, Baker GA, Havers S. Reality monitoring and psychotic hallucinations. *Br J Clin Psychol* 1991;30(Pt 3):213–22.



- 
- [177] Johns LC, Rossell S, Frith C, Ahmad F, Hemsley D, Kuipers E, McGuire PK. Verbal self-monitoring and auditory verbal hallucinations in patients with schizophrenia. *Psychol Med* 2001;31:705–15.
  - [178] Brebion G, Amador X, David A, Malaspina D, Sharif Z, Gorman JM. Positive symptomatology and source-monitoring failure in schizophrenia--an analysis of symptom-specific effects. *Psychiatry Res* 2000;95:119–31.
  - [179] Moritz S, Woodward TS, Ruff CC. Source monitoring and memory confidence in schizophrenia. *Psychol Med* 2003;33:131–9.
  - [180] Keefe RS, Arnold MC, Bayen UJ, Harvey PD. Source monitoring deficits in patients with schizophrenia ; a multinomial modelling analysis. *Psychol Med* 1999;29:903–14.
  - [181] Favrod J, Linder S, Jimenez X, Pomini V. Erreurs de mémoire et symptômes psychotiques : état des connaissances et perspectives thérapeutiques. In: Campanella S, Steel E, editors. *Psychopathologie et neurosciences questions actuelles de neurosciences cognitives et affectives*. Bruxelles: De Boeck Université; 2008. p. 123–47.
  - [182] Favrod J, Vianin P, Pomini V, Mast FW. A first step toward cognitive remediation of voices : A case study. *Cogn Behav Ther* 2006;35:159–63.
  - [183] American Psychiatric Association. Manuel diagnostique et statistique des troubles mentaux. 4<sup>e</sup> édition, texte révisé. Washington DC: American Psychiatric Association; 2000.
  - [184] Bell V, Halligan PW, Ellis HD. Explaining delusions: A cognitive perspective. *Trends Cogn Sci* 2006;10:219–26.
  - [185] Myin-Germeyns I, Nicolson NA, Delespaul PA. The context of delusional experiences in the daily life of patients with schizophrenia. *Psychol Med* 2001;31:489–98.
  - [186] Garety PA, Freeman D, Jolley S, Dunn G, Bebbington PE, Fowler DG, et al. Reasoning, emotions, and delusional conviction in psychosis. *J Abnorm Psychol* 2005;114:373–84.
  - [187] Kuhn T. The structure of scientific revolutions. Chicago (IL): University of Chicago Press; 1962.
  - [188] Bell V, Grech E, Maiden C, Halligan PW, Ellis HD. ‘Internet delusions’: A case series and theoretical integration. *Psychopathology* 2005;38:144–50.
  - [189] Bell V, Maiden C, Munoz-Solomando A, Reddy V. ‘Mind control’ experiences on the internet : Implications for the psychiatric diagnosis of delusions. *Psychopathology* 2006;39:87–91.
  - [190] Oltmanns TF. Approaches to the definition and study of delusions. In: Maher Tfoba, editor. *Delusional beliefs*. New York: Wiley; 1988. p. 3–12.
  - [191] Freeman D. Suspicious minds: The psychology of persecutory delusions. *Clin Psychol Rev* 2007;27:425–57.
  - [192] Stompe T, Ortwein-Swoboda G, Ritter K, Schanda H. Old wine in new bottles? Stability and plasticity of the contents of schizophrenic delusions. *Psychopathology* 2003;36:6–12.
  - [193] Bell V, Halligan PW, Ellis HD. Are anomalous perceptual experiences necessary for delusions ? *J Nerv Ment Dis* 2008;196:3–8.
  - [194] Freeman D, Gittins M, Pugh K, Antley A, Slater M, Dunn G. What makes one person paranoid and another person anxious? The differential prediction of social anxiety and persecutory ideation in an experimental situation. *Psychol Med* 2008;38:1121–32.
  - [195] Peters E, Garety P. Cognitive functioning in delusions: A longitudinal analysis. *Behav Res Ther* 2006;44:481–514.
  - [196] Van Dael F, Versmissen D, Janssen I, Myin-Germeyns I, van Os J, Krabbendam L. Data gathering: Biased in psychosis? *Schizophr Bull* 2006;32:341–51.
  - [197] Freeman D, Garety P, Kuipers E, Colbert S, Jolley S, Fowler D, et al. Delusions and decision-making style: Use of the need for closure scale. *Behav Res Ther* 2006;44:1147–58.

- [198] Colbert SM, Peters ER. Need for closure and jumping-to-conclusions in delusion-prone individuals. *J Nerv Ment Dis* 2002;190:27–31.
- [199] Moritz S, Woodward TS, Chen E. Investigation of metamemory dysfunctions in first-episode schizophrenia. *Schizophr Res* 2006;81:247–52.
- [200] Moritz S, Woodward TS. The contribution of metamemory deficits to schizophrenia. *J. Abnorm Psychol* 2006;115:15–25.
- [201] Jou J, Matus YE, Aldridge JW, Rogers DM, Zimmerman RL. How similar is false recognition to veridical recognition objectively and subjectively? *Mem Cognit* 2004;32:824–40.
- [202] Moritz S, Woodward TS, Whitman JC, Cuttler C. Confidence in errors as a possible basis for delusions in schizophrenia. *J Nerv Ment Dis* 2005;193:9–16.
- [203] Moritz S, Woodward TS. A generalized bias against disconfirmatory evidence in schizophrenia. *Psychiatry Res* 2006;142:157–65.
- [204] Woodward TS, Buchy L, Moritz S, Liotti M. A bias against disconfirmatory evidence is associated with delusion proneness in a nonclinical sample. *Schizophr Bull* 2007;33(4):1023–8.
- [205] Bentall RP, Corcoran R, Howard R, Blackwood N, Kinderman P. Persecutory delusions: A review and theoretical integration. *Clin Psychol Rev* 2001;21:1143–92.
- [206] Kinderman P, Bentall RP. Causal attributions in paranoia and depression: Internal, personal, and situational attributions for negative events. *J Abnorm Psychol* 1997;106:341–5.
- [207] Martin JA, Penn DL. Attributional style in schizophrenia: An investigation in outpatients with and without persecutory delusions. *Schizophr Bull* 2002;28:131–41.
- [208] McKay R, Langdon R, Coltheart M. Paranoia, persecutory delusions and attributional biases. *Psychiatry Res* 2005;136:233–45.
- [209] Randall F, Corcoran R, Day JC, Bentall RP. Attention, theory of mind, and causal attributions in people with persecutory delusions: A preliminary investigation. *Cognit Neuropsychiatry* 2003;8:287–94.
- [210] Frith CD. Schizophrenia and theory of mind. *Psychol Med* 2004;34:385–9.
- [211] Garety PA, Freeman D. Cognitive approaches to delusions : A critical review of theories and evidence. *Br J Clin Psychol* 1999;38(Pt 2):113–54.
- [212] Kelemen O, Erdelyi R, Pataki I, Benedek G, Janka Z, Keri S. Theory of mind and motion perception in schizophrenia. *Neuropsychology* 2005;19:494–500.
- [213] Mazza M, De Risio A, Surian L, Roncone R, Casacchia M. Selective impairments of the theory of mind in people with schizophrenia. *Schizophr Res* 2001;47:299–308.
- [214] Greig TC, Bryson GJ, Bell MD. Theory of mind performance in schizophrenia: Diagnostic, symptom, and neuropsychological correlates. *J Nerv Ment Dis* 2004;192:12–8.
- [215] Phillips ML, Senior C, David AS. Perception of threat in schizophrenics with persecutory delusions : An investigation using visual scan paths. *Psychol Med* 2000;30:157–67.
- [216] Freeman D, Garety P, Fowler D, Kuipers E, Dunn G, Bebbington P, Hadley C. The london-east anglia randomized controlled trial of cognitive-behaviour therapy for psychosis. IV: Self-esteem and persecutory delusions. *Br J Clin Psychol* 1998;37:415–30.
- [217] Bentall RP, Kinderman P, Kaney S. The self, attributional processes and abnormal beliefs: Towards a model of persecutory delusions. *Behav Res Ther* 1994;32:331–41.
- [218] Freeman D, Garety PA, Kuipers E, Fowler D, Bebbington PE. A cognitive model of persecutory delusions. *Br J Clin Psychol* 2002;41:331–47.
- [219] Drake RJ, Pickles A, Bentall RP, Kinderman P, Haddock G, Tarrier N, Lewis SW. The evolution of insight, paranoia and depression during early schizophrenia. *Psychol Med* 2004;34:285–92.
- [220] Rosse RB, Kendrick K, Wyatt RJ, Isaac A, Deutsch SI. Gaze discrimination in patients with schizophrenia: Preliminary report. *Am J Psychiatry* 1994;151:919–21.

- 
- [221] Franck N, Daprati E, Michel F, Saoud M, Dalery J, Marie-Cardine M, Georgieff N. Gaze discrimination is unimpaired in schizophrenia. *Psychiatry Res* 1998;81:67–75.
  - [222] Franck N, Montoute T, Labruyere N, Tiberghien G, Marie-Cardine M, Dalery J, et al. Gaze direction determination in schizophrenia. *Schizophr Res* 2002;56:225–34.
  - [223] Hooker C, Park S. You must be looking at me: The nature of gaze perception in schizophrenia patients. *Cognit Neuropsychiatry* 2005;10:327–45.
  - [224] Green MJ, Williams LM, Davidson D. Visual scanpaths and facial affect recognition in delusion-prone individuals : Increased sensitivity to threat? *Cognit Neuropsychiatry* 2003;8:19–41.
  - [225] Phillips ML, David AS. Visual scan paths are abnormal in deluded schizophrenics. *Neuropsychologia* 1997;35:99–105.
  - [226] Phillips ML, David AS. Abnormal visual scan paths: A psychophysiological marker of delusions in schizophrenia. *Schizophr Res* 1998;29:235–45.
  - [227] Williams LM, Loughland CM, Gordon E, Davidson D. Visual scanpaths in schizophrenia: Is there a deficit in face recognition? *Schizophr Res* 1999;40:189–99.
  - [228] Loughland CM, Williams LM, Gordon E. Visual scanpaths to positive and negative facial emotions in an outpatient schizophrenia sample. *Schizophr Res* 2002;55:159–70.
  - [229] Streit M, Wolwer W, Gaebel W. Facial-affect recognition and visual scanning behaviour in the course of schizophrenia. *Schizophr Res* 1997;24:311–7.
  - [230] Colbert SM, Peters ER. Need for closure and jumping-to-conclusions in delusion-prone individuals. *J Nerv Ment Dis* 2002;190:27–31.
  - [231] Freeman D, Garety PA, Kuipers E. Persecutory delusions : Developing the understanding of belief maintenance and emotional distress. *Psychol Med* 2001;31:1293–306.
  - [232] Freeman D, Garety PA, Kuipers E, Fowler D, Bebbington PE, Dunn G. Acting on persecutory delusions: The importance of safety seeking. *Behav Res Ther* 2007;45:89–99.
  - [233] Gaynor K, Ward T, Garety P, Peters E. The role of safety-seeking behaviours in maintaining threat appraisals in psychosis. *Behav Res Ther* 2013;51:75–81.
  - [234] Ward TA, Gaynor KJ, Hunter MD, Woodruff PW, Garety PA, Peters ER. Appraisals and responses to experimental symptom analogues in clinical and nonclinical individuals with psychotic experiences. *Schizophr Bull* 2014;40:845–55.
  - [235] Chaix J, Ma E, Nguyen A, Ortiz Collado MA, Rexhaj S, Favrod J. Safety-seeking behaviours and verbal auditory hallucinations in schizophrenia. *Psychiatry Res* 2014;.
  - [236] Stip E, Letourneau G. Psychotic symptoms as a continuum between normality and pathology. *Can J Psychiatry* 2009;54:140–51.
  - [237] Lund PM. Marathon volleyball: Changes after 61 hours play. *Br J Sports Med* 1985;19:228–9.
  - [238] Grassian S. Psychopathological effects of solitary confinement. *Am J Psychiatry* 1983;140:1450–4.
  - [239] Slade PD. Sensory deprivation and clinical psychiatry. *Br J Hosp Med* 1984;32:256–60.
  - [240] Brugger P, Regard M, Landis T, Oelz O. Hallucinatory experiences in extreme-altitude climbers. *Neuropsychiatry Neuropsychol Behav Neurol* 1999;12:67–71.
  - [241] Zimbardo PG, Andersen SM, Kabat LG. Induced hearing deficit generates experimental paranoia. *Science* 1981;212:1529–31.
  - [242] Nasar S. Un cerveau d'exception. De la schizophrénie au prix nobel, la vie singulière de John Forbes Nash. Paris: Calmann-Lévy; 2001.
  - [243] Kahane M. Nijinsky, 1889-1950. Paris: Éditions de la Réunion des musées nationaux; 2000.
  - [244] Scheder D. L'auto jaune. Parcours au travers d'une folie. Lausanne: Favre SA; 2005.
  - [245] Chadwick P. Schizophrenia: The positive perspective-explorations at the outer reaches of human experience. London: Taylor & Francis, Inc; 1997.

- [246] Lukoff D, Wallace CJ, Liberman RP, Burke K. A holistic program for chronic schizophrenic patients. *Schizophr Bull* 1986;12:274–82.
- [247] Nigl AJ, Jackson B. Electromyograph biofeedback as an adjunct to standard psychiatric treatment. *J Clin Psychiatry* 1979;40:433–6.
- [248] Durham RC, Guthrie M, Morton RV, Reid DA, Treiving LR, Fowler D, Macdonald RR. Tayside-five clinical trial of cognitive-behavioural therapy for medication-resistant psychotic symptoms. Results to 3-month follow-up. *Br J Psychiatry* 2003;182:303–11.
- [249] Pinto A, La Pia S, Mennella R, Giorgio D, DeSimone L. Cognitive-behavioral therapy and clozapine for clients with treatment-refractory schizophrenia. *Psychiatr Serv* 1999;50:901–4.
- [250] Hawkins RC 2nd, Doell SR, Lindseth P, Jeffers V, Skaggs S. Anxiety reduction in hospitalized schizophrenics through thermal biofeedback and relaxation training. *Percept Mot Skills* 1980;51:475–82.
- [251] Van Hassel JH, Bloom LJ, Gonzalez AM. Anxiety management with schizophrenic outpatients. *J Clin Psychol* 1982;38:280–5.
- [252] Haddock G, Bentall R, Slade PD. Psychological treatment of auditory hallucinations: Focusing or distraction? In: Haddock G, Slade PD, editors. *Cognitive-behavioural interventions with psychotic disorders*. London: Routledge; 1997. p. 45–70.
- [253] Haddock G, Slade PD, Bentall RP, Reid D, Faragher EB. A comparison of the long-term effectiveness of distraction and focusing in the treatment of auditory hallucinations. *Br J Med Psychol* 1998;71(Pt 3):339–49.
- [254] Chadwick P, Newman Taylor K, Abba N. Mindfulness groups for people with psychosis. *Behavioural and Cognitive Psychotherapy* 2005;33:351–9.
- [255] Segal ZV, Williams JMG, Teasdale JD. *La thérapie cognitive basée sur la pleine conscience pour la dépression*. Bruxelles: De Boeck; 2006.
- [256] Linehan M. *Traitement cognitivo-comportemental du trouble de personnalité éat-limite*. Genève: Médecine & Hygiène; 2000.
- [257] Garcia JM, Pérez M. Act as a treatment of psychotic symptoms. The case of auditory hallucinations. *Analisis y Modificacìon de Conducta* 2001;27:455–72.
- [258] Bach P, Hayes SC. The use of acceptance and commitment therapy to prevent the rehospitalization of psychotic patients: A randomized controlled trial. *J Consult Clin Psychol* 2002;70:1129–39.
- [259] Crawford-Walker CJ, King A, Chan S. Distraction techniques for schizophrenia. *Cochrane Database Syst Rev* 2005 : CD004717.
- [260] Morrison AP. Cognitive behaviour therapy for auditory hallucinations without concurrent medication: A singel case. *Behavioural and Cognitive Psychotherapy* 1994;22:259–64.
- [261] Morrison AP, Haddock G, Tarrier N. Intrusive thoughts and auditory hallucination: A cognitive approach. *Behavioural and Cognitive Psychotherapy* 1995;23:265–80.
- [262] Romme MA, Escher AD. The new approach: A dutch experiment. In: Romme MA, Escher AD, editors. *Accepting voices*. London: Mind; 1993.
- [263] Wegner DM, Schneider DJ, Carter SR 3rd, White TL. Paradoxical effects of thought suppression. *J Pers Soc Psychol* 1987;53:5–13.
- [264] Wegner DM. *The illusion of conscious will*. Cambridge (MA): MIT Press; 2003.
- [265] Hayes SC, Strosahl K, Wilson KG. *Acceptance and commitment therapy: An experiential approach to behavior change*. New York: Guilford Press; 1999.
- [266] Moritz S, Woodward TS, Metacognition Study Group, Favrod J. *Entraînement des habiletés métacognitives pour les personnes atteintes de schizophrénie (emc)*. Hamburg: VanHam Campus Verlag; 2007.
- [267] Aghotor J, Pfueller U, Moritz S, Weisbrod M, Roesch-Ely D. Metacognitive training for patients with schizophrenia (mct): Feasibility and preliminary evidence for its efficacy. *J Behav Ther Exp Psychiatry* 2010;41:207–11.

- 
- [268] Balzan RP, Delfabbro PH, Galletly CA, Woodward TS. Metacognitive training for patients with schizophrenia: Preliminary evidence for a targeted, single-module programme. *Aust N Z J Psychiatry* 2014;48(12):1126–36.
  - [269] Briki M, Monnin J, Haffen E, Sechter D, Favrod J, Netillard C, et al. Metacognitive training for schizophrenia: A multicentre randomised controlled trial. *Schizophr Res* 2014;157:99–106.
  - [270] Erawati E, Keliat BA, Helena N, Hamid A. The influence of metacognitive training on delusion severity and metacognitive ability in schizophrenia. *J Psychiatr Ment Health Nurs* 2014;21(9):841–7.
  - [271] Favrod J, Maire A, Bardy S, Pernier S, Bonsack C. Improving insight into delusions: A pilot study of metacognitive training for patients with schizophrenia. *J Adv Nurs* 2011;67:401–7.
  - [272] Favrod J, Rexhaj S, Bardy S, Ferrari P, Hayoz C, Moritz S, et al. Sustained antipsychotic effect of metacognitive training in psychosis: A randomized-controlled study. *Eur Psychiatry* 2014;29:275–81.
  - [273] Kumar D, Zia Ul Haq M, Dubey I, Dotivala KN, Veqar Siddiqui S, Prakash R, et al. Effect of meta-cognitive training in the reduction of positive symptoms in schizophrenia. *European Journal of Psychotherapy & Counselling* 2010;12:149–58.
  - [274] Kuokkanen R, Lappalainen R, Repo-Tiihonen E, Tiihonen J. Metacognitive group training for forensic and dangerous non-forensic patients with schizophrenia : A randomised controlled feasibility trial. *Crim Behav and Men Health* 2014;.
  - [275] Moritz S, Kerstan A, Veckenstedt R, Randjbar S, Vitzthum F, Schmidt C, et al. Further evidence for the efficacy of a metacognitive group training in schizophrenia. *Behav Res Ther* 2011;49:151–7.
  - [276] Moritz S, Veckenstedt R, Andreou C, Bohn F, Hottenrott B, Leighton L, et al. Sustained and “sleepier” effects of group metacognitive training for schizophrenia: A randomized clinical trial. *JAMA Psychiatry* 2014;71:1103–11.
  - [277] Moritz S, Veckenstedt R, Randjbar S, Vitzthum F, Woodward TS. Antipsychotic treatment beyond antipsychotics: Metacognitive intervention for schizophrenia patients improves delusional symptoms. *Psychol Med* 2011;41:1823–32.
  - [278] Moritz S, Woodward TS. Metacognitive training for schizophrenia patients (mct) : A pilot study of feasibility, treatment adherence and subjective efficacy. *German Journal of Psychiatry* 2007;10:69–78.
  - [279] Ross K, Freeman D, Dunn G, Garety P. A randomized experimental investigation of reasoning training for people with delusions. *Schizophr Bull* 2011;37(2):324–33.
  - [280] van Oosterhout B, Krabbendam L, de Boer K, Ferwerda J, van der Helm M, Stant AD, van der Gaag M. Metacognitive group training for schizophrenia spectrum patients with delusions: A randomized controlled trial. *Psychol Med* 2014;44:3025–35.
  - [281] Khazaal Y, Favrod J, Azoulay S, Pomini V. « Michael’s game »: entraînement au raisonnement par hypothèses chez des patients présentant des symptômes psychotiques. *Revue Francophone de Clinique Comportementale et Cognitive* 2008;13:1–7.
  - [282] Khazaal Y, Favrod J, Libbrecht J, Finot SC, Azoulay S, Benzakin L, et al. A card game for the treatment of delusional ideas: A naturalistic pilot trial. *BMC Psychiatry* 2006;6:48.
  - [283] Khazaal Y, Favrod J, Azoulay S, Finot SC, Bernabotto M, Raffard S, et al. Michael’s game,” a card game for the treatment of psychotic symptoms. *Patient Educ Couns* 2011;83(2):210–6.
  - [284] Ellett L, Freeman D, Garety PA. The psychological effect of an urban environment on individuals with persecutory delusions: The camberwell walk study. *Schizophr Res* 2008;99:77–84.
  - [285] Erhart SM, Marder SR, Carpenter WT. Treatment of schizophrenia negative symptoms: Future prospects. *Schizophr Bull* 2006;32:234–7.

- [286] Lecrubier Y, Quintin P, Bouhassira M, Perrin E, Lancrenon S. The treatment of negative symptoms and deficit states of chronic schizophrenia: Olanzapine compared to amisulpride and placebo in a 6-month double-blind controlled clinical trial. *Acta Psychiatr Scand* 2006;114:319–27.
- [287] Stahl SM, Buckley PF. Negative symptoms of schizophrenia: A problem that will not go away. *Acta Psychiatr Scand* 2007;115:4–11.
- [288] Alphas L. An industry perspective on the nimh consensus statement on negative symptoms. *Schizophr Bull* 2006;32:225–30.
- [289] Dickerson F, Boronow JJ, Ringel N, Parente F. Social functioning and neurocognitive deficits in outpatients with schizophrenia: A 2-year follow-up. *Schizophr Res* 1999;37:13–20.
- [290] Milev P, Ho BC, Arndt S, Andreasen NC. Predictive values of neurocognition and negative symptoms on functional outcome in schizophrenia: A longitudinal first-episode study with 7-year follow-up. *Am J Psychiatry* 2005;162:495–506.
- [291] Bow-Thomas CC, Velligan DI, Miller AL, Olsen J. Predicting quality of life from symptomatology in schizophrenia at exacerbation and stabilization. *Psychiatry Res* 1999;86:131–42.
- [292] Tarrier N, Wykes T. Is there evidence that cognitive behaviour therapy is an effective treatment for schizophrenia? A cautious or cautionary tale? *Behav Res Ther* 2004;42:1377–401.
- [293] Gould RA, Mueser KT, Bolton E, Mays V, Goff D. Cognitive therapy for psychosis in schizophrenia: An effect size analysis. *Schizophr Res* 2001;48:335–42.
- [294] Petersen L, Jeppesen P, Thorup A, Abel MB, Ohlenschlaeger J, Christensen TO, et al. A randomised multicentre trial of integrated versus standard treatment for patients with a first episode of psychotic illness. *BMJ* 2005;331:602.
- [295] Thorup A, Petersen L, Jeppesen P, Ohlenschlaeger J, Christensen T, Krarup G, et al. Integrated treatment ameliorates negative symptoms in first episode psychosis--results from the danish opus trial. *Schizophr Res* 2005;79:95–105.
- [296] Rector NA, Seeman MV, Segal ZV. Cognitive therapy for schizophrenia: A preliminary randomized controlled trial. *Schizophr Res* 2003;63:1–11.
- [297] Overall JE, Gorham G. The brief psychiatric rating scale. *Psychol Rep* 1962;10:799–812.
- [298] Kay SR, Fiszbein A, Opler LA. The positive and negative syndrome scale (panss) for schizophrenia. *Schizophr Bull* 1987;13:261–76.
- [299] Alphas LD, Summerfelt A, Lann H, Muller RJ. The negative symptom assessment: A new instrument to assess negative symptoms of schizophrenia. *Psychopharmacol Bull* 1989;25:159–63.
- [300] Kirkpatrick B, Buchanan RW, McKenney PD, Alphas LD, Carpenter WT Jr. The schedule for the deficit syndrome: An instrument for research in schizophrenia. *Psychiatry Res* 1989;30:119–23.
- [301] Favrod J, Ernst F, Giuliani F, Bonsack C. Validation française de l'échelle d'expérience temporelle du plaisir. *Encephale* 2009;35:241–8.
- [302] Blanchard JJ, Cohen AS. The structure of negative symptoms within schizophrenia : Implications for assessment. *Schizophr Bull* 2006;32:238–45.
- [303] Foussias G, Remington G. Negative symptoms in schizophrenia: Avolition and oc-cam's razor. *Schizophr Bull* 2010;36:359–69.
- [304] Strauss GP, Horan WP, Kirkpatrick B, Fischer BA, Keller WR, Miski P, et al. Deconstructing negative symptoms of schizophrenia: Avolition-apathy and diminished expression clusters predict clinical presentation and functional outcome. *J Psychiatr Res* 2013;47:783–90.
- [305] Gard DE, Germans Gard M, Kring AM, John OP. Anticipatory and consummatory components of the experience of pleasure : A scale development study. *Journal of Research in Personality* 2006;40:1086–102.



- 
- [306] Klein D. Depression and anhedonia. In: Clark DC, Fawcett J, editors. *Anhedonia and affect deficit states*. New York: PMA Publishing; 1984. p. 1–14.
  - [307] Kring AM. Emotion in schizohrenia: Old mystery, new understanding. *Current Directions in Psychological Science* 1999;8:160–3.
  - [308] Gard DE, Kring AM, Germans MK, Werner K. *Emotion in the daily life of patients with schizophrenia*. Boulder (CO): Society for Research in Psychopathology; 2000.
  - [309] Favrod J, Giuliani F, Ernst F, Bonsack C. Anticipatory pleasure skills training: A new intervention to reduce anhedonia in schizophrenia. *Perspect Psychiatr Care* 2010;46:171–81.
  - [310] Cassar R, Applegate E, Bentall RP. Poor savouring and low self-efficacy are predictors of anhedonia in patients with schizophrenia spectrum disorders. *Psychiatry Res* 2013;210:830–4.
  - [311] Quoidbach J, Berry EV, Hansenne M, Mikolajczak M. Positive emotion regulation and well-being: Comparing the impact of eight savoring and dampening strategies. *Personality and Individual Differences* 2010;49:368–73.
  - [312] Kring AM, Moran EK. Emotional response deficits in schizophrenia: Insights from affective science. *Schizophr Bull* 2008;34:819–34.
  - [313] Couture SM, Blanchard JJ, Bennett ME. Negative expectancy appraisals and defeatist performance beliefs and negative symptoms of schizophrenia. *Psychiatry Res* 2011;189:43–8.
  - [314] Grant PM, Beck AT. Defeatist beliefs as a mediator of cognitive impairment, negative symptoms, and functioning in schizophrenia. *Schizophr Bull* 2009;35:798–806.
  - [315] Kolb AY, Kolb DA. The learning way: Meta-cognitive aspects of experiential learning. *Simulation & Gaming* 2009;40:297–327.
  - [316] Horan WP, Green MF, Kring AM, Nuechterlein KH. Does anhedonia in schizophrenia reflect faulty memory for subjectively experienced emotions? *J Abnorm Psychol* 2006;115:496–508.
  - [317] Cungi C, Deglon C. *Cohérence cardiaque, nouvelles techniques pour faire face au stress*. Paris: Retz; 2009.
  - [318] Fioravanti M, Carlone O, Vitale B, Cinti ME, Clare L. A meta-analysis of cognitive deficits in adults with a diagnosis of schizophrenia. *Neuropsychol Rev* 2005;15:73–95.
  - [319] O’Carroll R. Cognitive impairment in schizophrenia. *Advances in Psychiatric Treatment* 2000;6:161–8.
  - [320] Pomini V, Neis L, Brenner HD, Hodel B, Roder V. *Thérapie psychologique des schizophrénies*. Hayen: Mardaga; 1998.
  - [321] Cochet A, Saoud M, Gabriele S, Broallier V, El Asmar C, Dalery J, D’Amato T. Impact de la remédiation cognitive dans la schizophrénie sur les stratégies de résolution de problèmes et l’autonomie sociale : Utilisation du logiciel rehacom. *Encephale* 2006;32:189–95.
  - [322] Vianin P. *Programme de remédiation cognitive pour patients présentant une schizophrénie ou un trouble associé*. Manuel du thérapeute. Charleroi: Socrate Editions Promarex; 2007.
  - [323] Spaulding WD, Reed D, Sullivan M, Richardson C, Weiler M. Effects of cognitive treatment in psychiatric rehabilitation. *Schizophr Bull* 1999;25:657–76.
  - [324] Chambon O, Marie-Cardine M. An evaluation of social skills training modules with schizophrenia inpatients in France. *International Review of Psychiatry* 1998;10:26–9.
  - [325] Chambon O, Marie-Cardine M, Cottraux J, Garcia JP, Favrod J, Deleu G. Impact d’un programme global d’entraînement aux habiletés sociales sur le fonctionnement social et la qualité de vie subjective de schizophrènes. *Journal de Thérapie Comportementale et Cognitive* 1995;37–43.
  - [326] Favrod J, Huguelet P, Chambon O. L’éducation au traitement neuroleptique peut-elle réduire les coûts ? Une étude pilote. *Encephale* 1996;22:331–6.

- [327] Stalberg G, Lichtenstein P, Sandin S, Hultman CM. Video-based assessment of interpersonal problem solving skills in patients with schizophrenia, their siblings and non-psychiatric controls. *Scand J Psychol* 2008;49:77–82.
- [328] Uçok A, Cakir S, Duman ZC, Discigil A, Kandemir P, Atli H. Cognitive predictors of skill acquisition on social problem solving in patients with schizophrenia. *Eur Arch Psychiatry Clin Neurosci* 2006;256:388–94.
- [329] Addington J, Addington D. Neurocognitive and social functioning in schizophrenia: A 2.5 year follow-up study. *Schizophr Res* 2000;44:47–56.
- [330] Favrod J, Caffaro M, Grossenbacher B, Rubio A, Von Turk A. Entraînement à la résolution de problèmes interpersonnels avec des personnes atteintes de schizophrénie dans différentes conditions de traitement. *Annales Médico-Psychologiques* 2000;158:302–11.
- [331] Granholm E, McQuaid JR, McClure FS, Link PC, Perivoliotis D, Gottlieb JD, et al. Randomized controlled trial of cognitive behavioral social skills training for older people with schizophrenia: 12-month follow-up. *J Clin Psychiatry* 2007;68:730–7.
- [332] Kern RS, Green MF, Mitchell S, Kopelowicz A, Mintz J, Liberman RP. Extensions of errorless learning for social problem-solving deficits in schizophrenia. *Am J Psychiatry* 2005;162:513–9.
- [333] Liberman RP, Glynn S, Blair KE, Ross D, Marder SR. In vivo amplified skills training: Promoting generalization of independent living skills for clients with schizophrenia. *Psychiatry* 2002;65:137–55.
- [334] Medalia A, Revheim N, Casey M. Remediation of problem-solving skills in schizophrenia : Evidence of a persistent effect. *Schizophr Res* 2002;57:165–71.
- [335] Donahoe CP, Carter MJ, Bloem WD, Hirsch GL, Laasi N, Wallace CJ. Assessment of interpersonal problem-solving skills. *Psychiatry* 1990;53:329–39.
- [336] Barbalat G, Chambon V, Franck N, Koechlin E, Farrer C. Organization of cognitive control within the lateral prefrontal cortex in schizophrenia. *Arch Gen Psychiatry* 2009;66:377–86.
- [337] Bazin N, Lefrere F, Passerieux C, Sarfati Y, Hardy-Bayle MC. Troubles formels de la pensée. Traduction française de l'échelle d'évaluation de la pensée, du langage et de la communication. *Encephale* 2002;28:109–19.
- [338] Bazin N, Sarfati Y, Lefrere F, Passerieux C, Hardy-Bayle MC. Scale for the evaluation of communication disorders in patients with schizophrenia: A validation study. *Schizophr Res* 2005;77:75–84.
- [339] Kerns JG, Berenbaum H. Cognitive impairments associated with formal thought disorder in people with schizophrenia. *J Abnorm Psychol* 2002;111:211–24.
- [340] Bora E, Yucel M, Pantelis C. Theory of mind impairment in schizophrenia: Meta-analysis. *Schizophr Res* 2009;109:1–9.
- [341] Sprong M, Schothorst P, Vos E, Hox J, van Engeland H. Theory of mind in schizophrenia : Meta-analysis. *Br J Psychiatry* 2007;191:5–13.
- [342] Shamay-Tsoory SG, Shur S, Barcai-Goodman L, Medlovich S, Harari H, Levkovitz Y. Dissociation of cognitive from affective components of theory of mind in schizophrenia. *Psychiatry Res* 2007;149:11–23.
- [343] Addington J, Penn D, Woods SW, Addington D, Perkins DO. Facial affect recognition in individuals at clinical high risk for psychosis. *Br J Psychiatry* 2008;192:67–8.
- [344] Roncone R, Falloon IR, Mazza M, De Risio A, Pollice R, Necozione S, et al. Is theory of mind in schizophrenia more strongly associated with clinical and social functioning than with neurocognitive deficits? *Psychopathology* 2002;35:280–8.
- [345] Bora E, Yucel M, Pantelis C. Theory of mind impairment: A distinct trait-marker for schizophrenia spectrum disorders and bipolar disorder? *Acta Psychiatr Scand* 2009;120(4):253–64.
- [346] Drury VM, Robinson EJ, Birchwood M. 'Theory of mind' skills during an acute episode of psychosis and following recovery. *Psychol Med* 1998;28:1101–12.

- 
- [347] Sarfati Y, Hardy-Bayle MC. How do people with schizophrenia explain the behaviour of others? A study of theory of mind and its relationship to thought and speech disorganization in schizophrenia. *Psychol Med* 1999;29:613–20.
  - [348] Baron-Cohen S, Leslie AM, Frith U. Does the autistic child have a “theory of mind”? *Cognition* 1985;21:37–46.
  - [349] Abu-Akel A, Abushua’leh K. ‘Theory of mind’ in violent and nonviolent patients with paranoid schizophrenia. *Schizophr Res* 2004;69:45–53.
  - [350] Mizrahi R, Korostil M, Starkstein SE, Zipursky RB, Kapur S. The effect of antipsychotic treatment on theory of mind. *Psychol Med* 2007;37:595–601.
  - [351] Bazin N, Brunet-Gouet E, Bourdet C, Kayser N, Falissard B, Hardy-Bayle MC, Passerieux C. Quantitative assessment of attribution of intentions to others in schizophrenia using an ecological video-based task: A comparison with manic and depressed patients. *Psychiatry Res* 2009;167:28–35.
  - [352] Madianos M, Economou M, Dafni O, Koukia E, Palli A, Rogakou E. Family disruption, economic hardship and psychological distress in schizophrenia: Can they be measured? *Eur Psychiatry* 2004;19:408–14.
  - [353] Magliano L, Fiorillo A, De Rosa C, Malangone C, Maj M. National Mental Health Project Working G. Family burden in long-term diseases: A comparative study in schizophrenia vs. Physical disorders. *Soc Sci Med* 2005;61:313–22.
  - [354] Moller-Leimkuhler AM. Burden of relatives and predictors of burden. Baseline results from the munich 5-year-follow-up study on relatives of first hospitalized patients with schizophrenia or depression. *Eur Arch Psychiatry Clin Neurosci* 2005;255:223–31.
  - [355] Moller-Leimkuhler AM. Multivariate prediction of relatives’ stress outcome one year after first hospitalization of schizophrenic and depressed patients. *Eur Arch Psychiatry Clin Neurosci* 2006;256:122–30.
  - [356] Hoenig J, Hamilton MW. The schizophrenic patient in the community and his effect on the household. *Int J Soc Psychiatry* 1966;12:165–76.
  - [357] Papastavrou E, Charalambous A, Tsangari H, Karayiannis G. The cost of caring: The relative with schizophrenia. *Scand J Caring Sci* 2010;24:817–23.
  - [358] Barrowclough C, Tarrier N, Johnston M. Distress, expressed emotion, and attributions in relatives of schizophrenia patients. *Schizophr Bull* 1996;22:691–702.
  - [359] Boye B, Bentsen H, Ulstein I, Notland TH, Lersbryggen A, Lingjaerde O, Malt UF. Relatives’ distress and patients’ symptoms and behaviours: A prospective study of patients with schizophrenia and their relatives. *Acta Psychiatr Scand* 2001;104:42–50.
  - [360] Mitsonis C, Voussoura E, Dimopoulos N, Psarra V, Kararizou E, Latzouraki E, et al. Patients associated with caregiver psychological distress in chronic schizophrenia. *Soc Psychiatry Psychiatr Epidemiol* 2012;47:331–7.
  - [361] Hodé Y. Le programme profamille: un outil pour modifier les relations soignants-soignés. *Confluences* 2007;17:40–2.
  - [362] Khazaal Y, Richard C, Preisig M, Zullino DF. Psychoéducation des proches de patients bipolaires: impact sur les patients. Rationnel, développements et perspectives. *Revue Médicale Suisse* 2004;504.
  - [363] Shimazu K, Shimodera S, Mino Y, Nishida A, Kamimura N, Sawada K, et al. Family psychoeducation for major depression: Randomised controlled trial. *Br J Psychiatry* 2011;198:385–90.
  - [364] Chien WT, Chan S, Morrissey J, Thompson D. Effectiveness of a mutual support group for families of patients with schizophrenia. *J Adv Nurs* 2005;51:595–608.
  - [365] Chien WT, Chan SW, Thompson DR. Effects of a mutual support group for families of chinese people with schizophrenia: 18-month follow-up. *Br J Psychiatry* 2006;189:41–9.
  - [366] Chien WT, Lee IY. The schizophrenia care management program for family caregivers of chinese patients with schizophrenia. *Psychiatr Serv* 2010;61:317–20.

- [367] Sellwood W, Wittkowski A, Tarrier N, Barrowclough C. Needs-based cognitive-behavioural family intervention for patients suffering from schizophrenia: 5-year follow-up of a randomized controlled effectiveness trial. *Acta Psychiatr Scand* 2007;116:447–52.
- [368] Koolae AK, Etemadi A. The outcome of family interventions for the mothers of schizophrenia patients in iran. *Int J Soc Psychiatry* 2010;56:634–46.
- [369] Sharif F, Shaygan M, Mani A. Effect of a psycho-educational intervention for family members on caregiver burdens and psychiatric symptoms in patients with schizophrenia in shiraz, iran. *BMC Psychiatry* 2012;12:48.
- [370] Kulhara P, Chakrabarti S, Avasthi A, Sharma A, Sharma S. Psychoeducational intervention for caregivers of indian patients with schizophrenia: A randomised-controlled trial. *Acta Psychiatr Scand* 2009;119:472–83.
- [371] Magliano L, Fiorillo A, Malangone C, De Rosa C, Maj M. Patient functioning and family burden in a controlled, real-world trial of family psychoeducation for schizophrenia. *Psychiatr Serv* 2006;57:1784–91.
- [372] Tas C, Danaci AE, Cubukcuoglu Z, Brune M. Impact of family involvement on social cognition training in clinically stable outpatients with schizophrenia - a randomized pilot study. *Psychiatry Res* 2012;195:32–8.
- [373] Motlova L, Dragomirecka E, Spaniel F, Goppoldova E, Zalesky R, Seleпова P, et al. Relapse prevention in schizophrenia: Does group family psychoeducation matter? One-year prospective follow-up field study. *Int J Psychiatry Clin Pract* 2006;10(1):38–44.
- [374] McWilliams S, Hill S, Mannion N, Fetherston A, Kinsella A, O'Callaghan E. Schizophrenia : A five-year follow-up of patient outcome following psycho-education for caregivers. *Eur Psychiatry* 2012;27:56–61.
- [375] Pharoah F, Mari J, Rathbone J, Wong W. Family intervention for schizophrenia. *Cochrane Database Syst Rev* 2010 : CD000088.
- [376] Gutierrez-Maldonado J, Caqueo-Urizar A. Effectiveness of a psycho-educational intervention for reducing burden in latin american families of patients with schizophrenia. *Qual Life Res* 2007;16:739–47.
- [377] Devaramane V, Pai NB, Vella SL. The effect of a brief family intervention on primary carer's functioning and their schizophrenic relatives levels of psychopathology in india. *Asian J Psychiatr* 2011;4:183–7.
- [378] Chien WT, Thompson DR, Norman I. Evaluation of a peer-led mutual support group for chinese families of people with schizophrenia. *Am J Community Psychol* 2008;42:122–34.
- [379] Magliano L, Orrico A, Fiorillo A, Del Vecchio H, Castiello G, Malangone C, et al. Family burden in bipolar disorders: Results from the italian mood disorders study (IMDS). *Epidemiol Psichiatri Soc* 2009;18:137–46.
- [380] Worakul P, Thavichachart N, Lueboonthavatchai P. Effects of psycho-educational program on knowledge and attitude upon schizophrenia of schizophrenic patients' caregivers. *J Med Assoc Thai* 2007;90:1199–204.
- [381] Gutierrez-Maldonado J, Caqueo-Urizar A, Ferrer-Garcia M. Effects of a psychoeducational intervention program on the attitudes and health perceptions of relatives of patients with schizophrenia. *Soc Psychiatry Psychiatr Epidemiol* 2009;44:343–8.
- [382] Sota S, Shimodera S, Kii M, Okamura K, Suto K, Suwaki M, et al. Effect of a family psychoeducational program on relatives of schizophrenia patients. *Psychiatry Clin Neurosci* 2008;62:379–85.
- [383] Duhamel F. La santé et la famille : une approche systémique en soins infirmiers. 2<sup>e</sup> édition Montréal (Québec): Édition de la Chenelière; 2007.
- [384] Cormier H, Guimond G, Joncas J, Leblanc G, Morin R, Vaillancourt S. Profamille, programme d'intervention de groupe auprès des familles de personnes atteintes de schizophrénie. Québec: Unité de psychiatrie sociale et préventive du CHUL et Centre de recherche de l'Université Laval Robert Giffard; 1991.

- 
- [385] Lecomte T, Leclerc C. Manuel de réadaptation psychiatrique. 2<sup>e</sup> édition Québec: Presses de l'Université du Québec; 2012.
  - [386] Feldmann R, Hornung WP, Prein B, Buchkremer G, Arolt V. Timing of psychoeducational psychotherapeutic interventions in schizophrenic patients. *Eur Arch Psychiatry Clin Neurosci* 2002;252:115–9.
  - [387] Duong L, Klitten LL, Moller RS, Ingason A, Jakobsen KD, Skjodt C, et al. Mutations in *nrxn1* in a family multiply affected with brain disorders: *Nrxn1* mutations and brain disorders. *Am J Med Genet B Neuropsychiatr Genet* 2012;159B:354–8.
  - [388] Rummel-Kluge C, Pitschel-Walz G, Bauml J, Kissling W. Psychoeducation in schizophrenia-results of a survey of all psychiatric institutions in germany, austria, and switzerland. *Schizophr Bull* 2006;32:765–75.
  - [389] Lincoln TM, Wilhelm K, Nestoriuc Y. Effectiveness of psychoeducation for relapse, symptoms, knowledge, adherence and functioning in psychotic disorders: A meta-analysis. *Schizophr Res* 2007;96:232–45.
  - [390] Rexhaj S, Python NV, Morin D, Bonsack C, Favrod J. Correlational study: Illness representations and coping styles in caregivers for individuals with schizophrenia. *Ann Gen Psychiatry* 2013;12:27.
  - [391] Chien WT, Thompson DR. An rct with three-year follow-up of peer support groups for chinese families of persons with schizophrenia. *Psychiatr Serv* 2013;64:997–1005.
  - [392] Bonsack C, Morandi S, Favrod J, Conus P. Le stigmate de la « folie » : de la fatalité au rétablissement. *Rev Med Suisse* 2013;9:588–92.
  - [393] Twamley EW, Narvaez JM, Becker DR, Bartels SJ, Jeste DV. Supported employment for middle-aged and older people with schizophrenia. *Am J Psychiatr Rehabil* 2008;11:76–89.
  - [394] Nuechterlein KH, Subotnik KL, Turner LR, Ventura J, Becker DR, Drake RE. Individual placement and support for individuals with recent-onset schizophrenia: Integrating supported education and supported employment. *Psychiatr Rehabil J* 2008;31:340–9.
  - [395] Cook JA, Leff HS, Blyler CR, Gold PB, Goldberg RW, Mueser KT, et al. Results of a multisite randomized trial of supported employment interventions for individuals with severe mental illness. *Arch Gen Psychiatry* 2005;62:505–12.
  - [396] Rummel-Kluge C, Stiegler-Kotzor M, Schwarz C, Hansen WP, Kissling W. Peer-counseling in schizophrenia: Patients consult patients. *Patient Educ Couns* 2008;70:357–62.
  - [397] West C. Powerful choices: Peer support and individualized medication self-determination. *Schizophr Bull* 2009;37(3):445–50.
  - [398] Favrod J, Nguyen A, Rexhaj S, Ferrari P, Ortiz A, Bonsack C. Back to the patient, l'informatique dans les soins. *Santé Mentale* 2014;74–7.
  - [399] Bonsack C, Favrod J. De la réhabilitation au rétablissement : l'expérience de lausanne. *L'Information Psychiatrique* 2013;89:227–32.
  - [400] Rogers JA, Vergare MJ, Baron RC, Salzer MS. Barriers to recovery and recommendations for change : The pennsylvania consensus conference on psychiatry's role. *Psychiatr Serv* 2007;58:1119–23.
  - [401] Latimer E. Economic considerations associated with assertive community treatment and supported employment for people with severe mental illness. *J Psychiatry Neurosci* 2005;30:355–9.
  - [402] Liberman RP. Dissemination and adoption of social skills training: Social validation of an evidence-based treatment for the mentally disabled. *J Ment Health* 2007;16:595–623.
  - [403] Tarrier N. Cognitive behaviour therapy for schizophrenia - a review of development, evidence and implementation. *Psychother Psychosom* 2005;74:136–44.